

www.BurdetteDental.com

Dr.'s Name		Phone #	
Address			
Date Wanted	Patient		
	☐ MALE		AGE





\$109 MicroZ° Layered Zirconia

*Price per unit upto three units


\$109 Per Unit on **MicroZ**° Layered Zirconia Expires December 31, 2018

Signature	License #	Date

NEED SUPPLIES?	ENCLOSED WITH CASE	SHADE INSTRUCTIONS
☐ BOXES ☐ BIO BAGS	☐ IMPRESSION ☐ BITE	SHADE NO
☐ FEDEX SHIPPING LABELS ☐ RX'S	☐ MODELS ☐ PHOTOS	
OTHER	OTHER	

FOR LAB USE ONLY		
Contact / LMOM:	Date:	
RE:		
Result:		
	Initial:	

Ideally, the preparation includes a circumferential shoulder or chamfer with a horizontal angle of at least 5°. The vertical preparation angle should be at least 4°. The inside angle of the shoulder preparation must be given a rounded contour. All occlusal and incisal edges should also be rounded. The marginal edge of the preparation needs to be continuous and clearly visible. A bevel should be avoided. For posterior and anterior teeth, a supragingival margin poses no problems. Due to the tooth-colored framework, very esthetic results can be achieved.



Do:

- Change nothing from your usual crown and bridge / P.F.M. technique. Prep, imp, bite, cement.
- As with any ceramic restoration, proper reduction of tooth/prep helps us build a restoration of unequalled esthetics.

Don't:

- Leave sharp tips on corners of the prep.
- Do a shoulder prep. This is not necessary for Crystal or Diamond crowns. If you do prep a shoulder, please do
 not prep a trough on the shoulder (the scanner does not read these well).
- Have patient force crown to place with cotton roll or anything else.
- Internally relieve all ceramic crown to seat. This will cause micro-fractures and in return will cause breakage in the future. If you feel relief needs to be done for additional seating, relief must be done to prep.

TERMS & POLICIES



www.BurdetteDental.com

Dr.'s Name		Phone #	
Address			
Date Wanted	Patient		
	☐ MALE		AGE





\$109 MicroZ° Layered Zirconia

*Price per unit upto three units


\$109 Per Unit on **MicroZ**° Layered Zirconia Expires December 31, 2018

Signature	License #	Date

NEED SUPPLIES?	ENCLOSED WITH CASE	SHADE INSTRUCTIONS
☐ BOXES ☐ BIO BAGS	☐ IMPRESSION ☐ BITE	SHADE NO
☐ FEDEX SHIPPING LABELS ☐ RX'S	☐ MODELS ☐ PHOTOS	
OTHER	OTHER	

FOR LAB USE ONLY		
Contact / LMOM:	Date:	
RE:		
Result:		
	Initial:	

Ideally, the preparation includes a circumferential shoulder or chamfer with a horizontal angle of at least 5°. The vertical preparation angle should be at least 4°. The inside angle of the shoulder preparation must be given a rounded contour. All occlusal and incisal edges should also be rounded. The marginal edge of the preparation needs to be continuous and clearly visible. A bevel should be avoided. For posterior and anterior teeth, a supragingival margin poses no problems. Due to the tooth-colored framework, very esthetic results can be achieved.



Do:

- Change nothing from your usual crown and bridge / P.F.M. technique. Prep, imp, bite, cement.
- As with any ceramic restoration, proper reduction of tooth/prep helps us build a restoration of unequalled esthetics.

Don't:

- Leave sharp tips on corners of the prep.
- Do a shoulder prep. This is not necessary for Crystal or Diamond crowns. If you do prep a shoulder, please do
 not prep a trough on the shoulder (the scanner does not read these well).
- Have patient force crown to place with cotton roll or anything else.
- Internally relieve all ceramic crown to seat. This will cause micro-fractures and in return will cause breakage in the future. If you feel relief needs to be done for additional seating, relief must be done to prep.

TERMS & POLICIES



www.BurdetteDental.com

Dr.'s Name		Phone #	
Address			
Date Wanted	Patient		
	☐ MALE		AGE





\$109 MicroZ° Layered Zirconia

*Price per unit upto three units


\$109 Per Unit on **MicroZ**° Layered Zirconia Expires December 31, 2018

Signature	License #	Date

NEED SUPPLIES?	ENCLOSED WITH CASE	SHADE INSTRUCTIONS
☐ BOXES ☐ BIO BAGS	☐ IMPRESSION ☐ BITE	SHADE NO
☐ FEDEX SHIPPING LABELS ☐ RX'S	☐ MODELS ☐ PHOTOS	
OTHER	OTHER	

FOR LAB USE ONLY		
Contact / LMOM:	Date:	
RE:		
Result:		
	Initial:	

Ideally, the preparation includes a circumferential shoulder or chamfer with a horizontal angle of at least 5°. The vertical preparation angle should be at least 4°. The inside angle of the shoulder preparation must be given a rounded contour. All occlusal and incisal edges should also be rounded. The marginal edge of the preparation needs to be continuous and clearly visible. A bevel should be avoided. For posterior and anterior teeth, a supragingival margin poses no problems. Due to the tooth-colored framework, very esthetic results can be achieved.



Do:

- Change nothing from your usual crown and bridge / P.F.M. technique. Prep, imp, bite, cement.
- As with any ceramic restoration, proper reduction of tooth/prep helps us build a restoration of unequalled esthetics.

Don't:

- Leave sharp tips on corners of the prep.
- Do a shoulder prep. This is not necessary for Crystal or Diamond crowns. If you do prep a shoulder, please do
 not prep a trough on the shoulder (the scanner does not read these well).
- Have patient force crown to place with cotton roll or anything else.
- Internally relieve all ceramic crown to seat. This will cause micro-fractures and in return will cause breakage in the future. If you feel relief needs to be done for additional seating, relief must be done to prep.

TERMS & POLICIES



www.BurdetteDental.com

Dr.'s Name		Phone #	
Address			
Date Wanted	Patient		
	☐ MALE		AGE





\$109 MicroZ° Layered Zirconia

*Price per unit upto three units


\$109 Per Unit on **MicroZ**° Layered Zirconia Expires December 31, 2018

Signature	License #	Date

NEED SUPPLIES?	ENCLOSED WITH CASE	SHADE INSTRUCTIONS
☐ BOXES ☐ BIO BAGS	☐ IMPRESSION ☐ BITE	SHADE NO
☐ FEDEX SHIPPING LABELS ☐ RX'S	☐ MODELS ☐ PHOTOS	
OTHER	OTHER	

FOR LAB USE ONLY		
Contact / LMOM:	Date:	
RE:		
Result:		
	Initial:	

Ideally, the preparation includes a circumferential shoulder or chamfer with a horizontal angle of at least 5°. The vertical preparation angle should be at least 4°. The inside angle of the shoulder preparation must be given a rounded contour. All occlusal and incisal edges should also be rounded. The marginal edge of the preparation needs to be continuous and clearly visible. A bevel should be avoided. For posterior and anterior teeth, a supragingival margin poses no problems. Due to the tooth-colored framework, very esthetic results can be achieved.



Do:

- Change nothing from your usual crown and bridge / P.F.M. technique. Prep, imp, bite, cement.
- As with any ceramic restoration, proper reduction of tooth/prep helps us build a restoration of unequalled esthetics.

Don't:

- Leave sharp tips on corners of the prep.
- Do a shoulder prep. This is not necessary for Crystal or Diamond crowns. If you do prep a shoulder, please do
 not prep a trough on the shoulder (the scanner does not read these well).
- Have patient force crown to place with cotton roll or anything else.
- Internally relieve all ceramic crown to seat. This will cause micro-fractures and in return will cause breakage in the future. If you feel relief needs to be done for additional seating, relief must be done to prep.

TERMS & POLICIES



www.BurdetteDental.com

Dr.'s Name		Phone #	
Address			
Date Wanted	Patient		
	☐ MALE		AGE





\$109 MicroZ° Layered Zirconia

*Price per unit upto three units


\$109 Per Unit on **MicroZ**° Layered Zirconia Expires December 31, 2018

Signature	License #	Date

NEED SUPPLIES?	ENCLOSED WITH CASE	SHADE INSTRUCTIONS
☐ BOXES ☐ BIO BAGS	☐ IMPRESSION ☐ BITE	SHADE NO
☐ FEDEX SHIPPING LABELS ☐ RX'S	☐ MODELS ☐ PHOTOS	
OTHER	OTHER	

FOR LAB USE ONLY		
Contact / LMOM:	Date:	
RE:		
Result:		
	Initial:	

Ideally, the preparation includes a circumferential shoulder or chamfer with a horizontal angle of at least 5°. The vertical preparation angle should be at least 4°. The inside angle of the shoulder preparation must be given a rounded contour. All occlusal and incisal edges should also be rounded. The marginal edge of the preparation needs to be continuous and clearly visible. A bevel should be avoided. For posterior and anterior teeth, a supragingival margin poses no problems. Due to the tooth-colored framework, very esthetic results can be achieved.



Do:

- Change nothing from your usual crown and bridge / P.F.M. technique. Prep, imp, bite, cement.
- As with any ceramic restoration, proper reduction of tooth/prep helps us build a restoration of unequalled esthetics.

Don't:

- Leave sharp tips on corners of the prep.
- Do a shoulder prep. This is not necessary for Crystal or Diamond crowns. If you do prep a shoulder, please do
 not prep a trough on the shoulder (the scanner does not read these well).
- Have patient force crown to place with cotton roll or anything else.
- Internally relieve all ceramic crown to seat. This will cause micro-fractures and in return will cause breakage in the future. If you feel relief needs to be done for additional seating, relief must be done to prep.

TERMS & POLICIES



www.BurdetteDental.com

Dr.'s Name		Phone #	
Address			
Date Wanted	Patient		
	☐ MALE		AGE





\$109 MicroZ° Layered Zirconia

*Price per unit upto three units


\$109 Per Unit on **MicroZ**° Layered Zirconia Expires December 31, 2018

Signature	License #	Date

NEED SUPPLIES?	ENCLOSED WITH CASE	SHADE INSTRUCTIONS
☐ BOXES ☐ BIO BAGS	☐ IMPRESSION ☐ BITE	SHADE NO
☐ FEDEX SHIPPING LABELS ☐ RX'S	☐ MODELS ☐ PHOTOS	
OTHER	OTHER	

FOR LAB USE ONLY		
Contact / LMOM:	Date:	
RE:		
Result:		
	Initial:	

Ideally, the preparation includes a circumferential shoulder or chamfer with a horizontal angle of at least 5°. The vertical preparation angle should be at least 4°. The inside angle of the shoulder preparation must be given a rounded contour. All occlusal and incisal edges should also be rounded. The marginal edge of the preparation needs to be continuous and clearly visible. A bevel should be avoided. For posterior and anterior teeth, a supragingival margin poses no problems. Due to the tooth-colored framework, very esthetic results can be achieved.



Do:

- Change nothing from your usual crown and bridge / P.F.M. technique. Prep, imp, bite, cement.
- As with any ceramic restoration, proper reduction of tooth/prep helps us build a restoration of unequalled esthetics.

Don't:

- Leave sharp tips on corners of the prep.
- Do a shoulder prep. This is not necessary for Crystal or Diamond crowns. If you do prep a shoulder, please do
 not prep a trough on the shoulder (the scanner does not read these well).
- Have patient force crown to place with cotton roll or anything else.
- Internally relieve all ceramic crown to seat. This will cause micro-fractures and in return will cause breakage in the future. If you feel relief needs to be done for additional seating, relief must be done to prep.

TERMS & POLICIES